NEWTON'S SECOND LAW OF MOTION— FORCE AND ACCELERATION

Objectives

- State the relationship between acceleration and net force. (6.1)
- State the relationship between acceleration and mass. (6.2)
- State and explain Newton's second law of motion. (6.3)
- List the factors that affect the force of friction between surfaces. (6.4)
- Distinguish between force and pressure. (6.5)
- Explain why the acceleration of an object in free fall does not depend upon the mass of the object. (6.6)
- List the factors that affect the air resistance force on an object. (6.7)

discover!

MATERIALS stopwatch, coffee filters, meter stick

EXPECTED OUTCOME The time of fall for all four trials should be the same. The results illustrate how the terminal velocity is proportional to the square root of the mass of the object divided by the time.

ANALYZE AND CONCLUDE

- The coffee filter accelerates at first and then moves at a constant velocity. The time is the same in all four trials.
 The same amount of time
- as the other trials 3. The mass of the object
- Ine mass of the obje affects its speed.

NEWTON'S SECOND LAW OF **NOTION – FORCE AND ACCELERATION**



n Chapter 2, we discussed the concept of mechanical equilibrium, $\Sigma F = 0$, which means that forces are balanced. In Chapter 3, we extended this idea to the law of inertia, again with balanced forces. In this chapter we consider what happens when forces aren't balanced when the net force is not zero—when an object is not zero when an object is not zero, and the ball accelerates. Its path through the air is not a straight line but curves downward due to gravity—again an acceleration. Most of the motion we see undergoes change. This chapter covers changes in motion—accelerated motion.

We learned that acceleration describes how quickly velocity changes. Specifically, it is the change in velocity per unit of time. Recall the definition of acceleration:

 $acceleration = \frac{change in velocity}{time interval}$

We will now focus on the cause of acceleration: force

discover

What Effect Does Air Resistance Have on Falling Objects?

- Use a stopwatch to determine the time required for a single coffee filter to fall one meter.
- Determine the time required for four coffee filters nested inside one another to fall two meters.
- **3.** Determine the time required for nine nested filters to fall a distance of three meters.
- If possible, measure the time of fall for sixteen nested filters dropped from a height of four meters.

Analyze and Conclude

- **1. Observing** What did you observe about the motion of a single filter as it fell? Did it appear to accelerate or did it move with a constant velocity? How did the time of fall compare for each of the four trials?
- Predicting How long do you think it would take for twenty-five nested coffee filters to fall through a distance of five meters?
- 3. Making Generalizations What determines the speed of similarly shaped objects falling under the influence of gravity and air resistance?

Force Causes Acceleration

accelerate. by striking the puck again, and the puck's motion changes again. to it) and the puck experiences a change in motion—it accelerates. puck is in equilibrium. Hit the puck (that is, apply an unbalanced force ice. The forces on it (gravity and the support force) are balanced, so the Consider an object at rest, such as a hockey puck on perfectly smooth 𝕑 Unbalanced forces acting on an object cause the object to forces and the puck moves at constant velocity. Apply another force When the hockey stick is no longer pushing it, there are no unbalanced

acceleration of an object, you must increase the net force acting on object. For example, after the boy kicks the football in Figure 6.1, is directly proportional to the net force acting on it. We write triple the force, its acceleration triples. We say an object's acceleration is the net force. Acceleration depends on the net force. To increase the previous chapter that the combination of forces acting on an object both gravity and air resistance act on the football. Recall from the it. Double the force on an object and its acceleration doubles. If you Most often, the force we apply is not the only force acting on an

acceleration \sim net force

The symbol \sim stands for "is directly proportional to."

CHECK: What causes an object to accelerate?

6.2 Mass Resists Acceleration

twice as much mass results in only half the acceleration. For three times will result in a decrease in the acceleration. The same force applied to the mass being pushed. **Tor a constant force, an increase in the mass** will accelerate much less than the empty cart. Acceleration depends on This relationship can be written as an equation: force, the acceleration produced is *inversely proportional* to the mass. the mass, one-third the acceleration results. In other words, for a given loaded shopping cart, as shown in Figure 6.2. The loaded shopping cart Push on an empty shopping cart. Then push equally hard on a heavily

acceleration $\sim \frac{1}{\text{mass}}$

quantity decreases by the same factor. Mathematically we see that as the denominator increases, the whole **Inversely** means that the two values change in opposite directions.

CHECK How does an increase in mass affect acceleration?



a straight line. remains at rest nor moves in FIGURE 6.1 Kick a football and it neither



FIGURE 6.2

is pushed depends on the mass that The acceleration produced

6.1 Force Causes Acceleration

evidence of zero net force. acceleration. Zero acceleration is Zero net force means zero act on the object are balanced. motion because the forces that there is no change in the state of the state of rest and the state of acceleration. Stress the idea that at constant velocity has zero students that an object traveling constant velocity. In both cases, zero acceleration applies both to Teaching Tip Remind

plane would speed up; if it were air resistance that acts on the and the thrust of the engines flying at a constant 900 km/h announces that the plane is more, the plane would slow force. If resistance were less, the 80,000 N, to produce a zero net plane's outside surface? What is the combined force of Zero, because velocity is constant. the acceleration of the airplane? is a constant 80,000 N. What is 诸 Ask Suppose a pilot

illustrate the previous situation body diagram on the board to Teaching Tip Draw a freedown.



cause the object to accelerate. CHECK acting on an object

Teaching Resources

- Reading and Study
- PresentationEXPRESS Workbook
- Conceptual Physics Alive!
- DVDs Newton's Second Law



CHAPTER 6

NEWTON'S SECOND LAW OF MOTION—FORCE AND ACCELERATION 87

6.2 Mass Resists Acceleration

inversely **Key Term**

although the acceleration of an object depends on the net force, Teaching Tip Point out that it also depends on the mass of

the object.

the acceleration. mass will result in a decrease in **CHECK** an increase in the **CONCEPT** For a constant force,

Teaching Resources

- Reading and Study Workbook
- PresentationEXPRESS
- Next-Time Question 6-1 Interactive Textbook

6.3 Newton's Second Law

Newton's second law Key Term

Teaching Tip Review the concept of inertia and discuss its role in Newton's second law.

discover!

MATERIALS spool of thread

same direction as the net force. of thread will accelerate in the EXPECTED OUTCOME The spool

- The spool will accelerate to the right.
- The spool will accelerate to the right.
- ω Yes. According to Newton's and an object's acceleration are always in the same direction. second law, the net force





6.3 Newton's Second Law

of the net force, is in the same direction as the net force, and is a net force on an object is directly proportional to the magnitude mass, an object's acceleration, and the net force on an object. but also on the object's mass. He came up with one of the most we move something depends not only on how hard we push or pull, inversely proportional to the mass of the object. 𝒞 Newton's second law states that the acceleration produced by Newton's second law describes the relationship among an object's important rules of nature ever proposed, his second law of motion Newton was the first to realize that the acceleration produced when

This relationship can be written as an equation:

acceleration
$$\sim \frac{\text{net force}}{\text{mass}}$$

acceleration, we get the exact equation kilograms (kg) for mass, and meters per second squared (m/s^2) for By using consistent units, such as newtons (N) for force

acceleration =
$$\frac{\text{net force}}{\text{mass}}$$

In briefest form, where *a* is acceleration, *F* is net force, and *m* is mass,

$$a = \frac{F}{m}$$

an object doubles its acceleration. Suppose instead that the mass is the mass are doubled, the acceleration will be unchanged doubled. Then acceleration will be halved. If both the net force and From this relationship we see that doubling the net force acting on The acceleration is equal to the net force divided by the mass.

CHECK: object's acceleration, and the net force on an object? **CONCEPT**: What is the relationship among an object's mass, an

discover!

Acceleration, Which Way?

- 1. Pull a spool of thread horizontally to the right by the spool. Which direction does the spool roll? the thread. The thread should be at the bottom of
- 2. Repeat step one with the thread at the top of the spool. Which direction does the spool roll?
- 3. Are the net force on an object and an object's acceleration always in the same direction? Why?





mass?

Answer: 6.3

can it attain if it is towing another car of equal

2 m/s², what acceleration If a car can accelerate at thinks

FIGURE 6.3

to its ability to produce large forces. The great acceleration of the racing car is due

do the math!

the car's acceleration. acceleration produced by a force of 2000 N? A car has a mass of 1000 kg. What is the You can use Newton's second law to solve for

a || т Е 1000 kg 2000 N 2000 kg·m/s² 1000 kg || 2 m/s²

If the force is 4000 N, what is the acceleration?

$$a = \frac{F}{m} = \frac{4000 \text{ N}}{1000 \text{ kg}} = \frac{4000 \text{ kg} \cdot \text{m/s}^2}{1000 \text{ kg}} = 4 \text{ m/s}^2$$

doubles the acceleration. Doubling the force on the same mass simply

and Solve problems go a step or two further for Then problem solving will be more meaningful. tant skill in physics. But first, learn the concepts more challenge. Solving problems is an importers familiarize you with equations, and the Think and Chug problems at the ends of many chaprelationships of basic physics concepts. The Plug size equations as guides to thinking about the cated problems in this book. Instead we emphathan these. We don't focus on solving compli-Physics problems are often more complicated

> of 1.5 m/s²? jet plane develop to achieve an acceleration How much force, or thrust, must a 30,000-kg

Newton's second law to read one meter per second squared. You can arrange give a mass of one kilogram an acceleration of tons (N). One newton is the force needed to (m/s²), then the force will be expressed in new-(kg) and its acceleration in meters per second If you know the mass of an object in kilograms

force = mass × acceleration

F = ma

= (30,000 kg)(1.5 m/s²) $= 45,000 \text{ kg} \cdot \text{m/s}^2$

= 45,000 N

units are multiplied together. The dot between kg and m/s² means that the



mass. Compare the resulting Apply equal forces to a masses. accelerations of the two large mass and a small

do the math!

problems are in Appendix F, example is Problem 52 in than numerical values. An solving involves numbers, a for more advanced students. Chapter 7 Assess. More such involves only symbols rather numbers, where the math problem is posed without problems. A kind of advanced who have done math-based familiar format to all students are straightforward. Problem The problems presented here

direction as the net force, and is of the net force, is in the same proportional to the magnitude acceleration produced by a net mass of the object. force on an object is directly inversely proportional to the CHECK law states that the

Teaching Resources

- Concept-Development Practice Book 6-1, 6-2,
- Problem-Solving Exercises in 6-3, 6-4
- Physics 4-1
- Laboratory Manual 18, 19, 20
- Probeware Lab Manual 4, 5
- PresentationEXPRESS
- Interactive Textbook
- Next-Time Question 6-2
- Conceptual Physics Alive!
- DVDs

Newton's Second Law

CHAPTER 6

6.4 Friction

Key Terms fluid, air resistance, free-body diagram

Teaching Tip Caution: We say that friction acts in a direction to oppose motion. But when you walk, the friction that acts on your shoes is in the same direction as your motion. But note that the contact of your foot on the floor is backward. Hence an opposite force of friction pushes you forward. Shoes push backward on the floor while the floor pushes your shoes and you forward. This will be clearer when Newton's third

CONCEPT: The force of friction **CHECK**: between the surfaces depends on the kinds of materials in contact and how much the surfaces are pressed together.

Teaching Resources

- Reading and Study Workbook
- Concept-Development Practice Book 6-5, 6-6
- Problem-Solving Exercises
- in Physics 4-2

 PresentationEXPRESS
- Interactive Textbook



FIGURE 6.4

A concrete road divider has a better design than a steel road divider for slowing an out-of-control, sideswiping car.

think!-

law is discussed.

Two forces act on a book resting on a table: its weight and the support force from the table. Does a force of friction act as well? Answer: 6.4

6.4 Friction

Friction is a force like any other force and affects motion. Friction acts on materials that are in contact with each other, and it always acts in a direction to oppose relative motion. When two solid objects come into contact, the friction is mainly due to irregularities in the two surfaces. When one object slides against another, it must either rise over the irregular bumps or else scrape them off. Either way requires force.

✓ The force of friction between the surfaces depends on the kinds of material in contact and how much the surfaces are pressed together. For example, rubber against concrete produces more friction than steel against steel. That's why concrete road dividers have replaced steel rails. The friction produced by a tire rubbing against the concrete is more effective in slowing the car than the friction produced by a steel car body sliding against a steel rail. Notice in Figure 6.4 that the concrete divider is wider at the bottom to ensure that the tire of a sideswiping car will make contact with the divider before the steel car body does.

Friction is not restricted to solids sliding or tending to slide over one another. Friction also occurs in liquids and gases. Both liquids and gases are called **fluids** because they flow. Fluid friction occurs as an object pushes aside the fluid it is moving through. Have you ever tried running a 100-m dash through waist-deep water? The friction of liquids is appreciable, even at low speeds. **Air resistance** is the friction acting on something moving through air. Air resistance is a very common form of fluid friction. You usually don't notice air resistance when walking or jogging, but you do notice it at the higher speeds that occur when riding a bicycle or skiing downhill.



Automobile Design The first automobiles were little more than horse carriages with engines. Over time, engineers came to realize that by reducing the frontal surface of cars and eliminating parts that stick out, the air resistance force on a car could be reduced. When a car cruises at a constant speed, the net force on the car is zero. By lowering the air resistance force at any speed, the amount of force needed by the engine is reduced, meaning better fuel economy. Over the years, cars have gotten sleeker, with teardrop-shaped bodies, and teardrop shapes around side mirrors. Door handles are set into the doors. Even wheel wells and the undersides of cars have been smoothed. Automotive engineers use computers to design cars with less air resistance and use wind tunnels to measure the cars' air resistance.



FIGURE 6.5 The direction of the force of friction always opposes the direction of motion. a. Push the crate to the right and friction acts toward the left. b. The sack falls downward and air fric-

tion acts upward.

When friction is present, an object may move with a constant velocity even when an outside force is applied to it. In such a case, the friction force just balances the applied force. The net force is zero, so there is no acceleration. For example, in Figure 6.5 the crate moves with a constant velocity when the force pushing it just balances the force of friction. The sack will also fall with a constant velocity once the force due to air resistance balances the sack's weight. A diagram showing all the forces acting on an object is called a **free-body diagram**.

CONCEPT: What factors affect the force of friction **CHECK** : between surfaces?

6.5 Applying Force—Pressure

Look at Figure 6.6. No matter how you place a book on a table, the force of the book on the table is the same. You can check this by placing a book in any position on a bathroom scale. You'll read the same weight in all cases. Balance a book in different positions on the palm of your hand. Although the force is always the same, you'll notice differences in the way the book presses against your palm. These differences are due to differences in the area of contact for each case. **©** For a constant force, an increase in the area of contact for per unit of area is called **pressure.** More precisely, when the force is perpendicular to the surface area,

$$pressure = \frac{force}{area of application}$$

In equation form,

$$=$$

 A F

Ρ

where *P* is the pressure and *A* is the area over which the force acts. Force, which is measured in newtons, is different from pressure. Pressure is measured in newtons per square meter, or **pascals** (Pa). One newton per square meter is equal to one pascal.



FIGURE 6.6

The upright book exerts the same force, but greater pressure, against the supporting surface.

6.5 Applying Force— Pressure

Key Terms pressure, pascal

Common Misconception

Force and pressure are the same. FACT Force is a push or a pull, while pressure is the amount of force per unit area.

a person stands on it with one a person stands on it with one pressure on something? Increase are two ways to increase the of contact has decreased. What foot in the air? Yes. The area against the scale increase when so the reading on the scale does weight does not change when foot in the air? No. The person's bathroom scale change when contact. the force; decrease the area of not change. Does the pressure the person stands on one foot, Ask Does the reading on a

Demonstration

Using a large demonstration spring scale, drag a block, wide side down, across a table. Repeat with the narrow side down. Show that the force required to pull the block across the table at constant velocity is independent of which surface is against the table. Explain that in both cases the weight of the block against the table is the same, but the *distribution of weight* is different.

Teaching Tip Define pressure. Give examples of different pressures: sharp points vs. blunt surfaces, a book resting on its cover vs. resting on its spine, etc.

In a recent repeat of the Bed of Nails demo (one we've done for many years) I had to swing harder than before to break the block. Why? Because the newer blocks are of more sturdy construction following the California earthquake of 1989. It is important that the block break, for otherwise, the person on the bed of nails gets more of the hammer's kinetic energy!

TAUL

Teaching Tip Note

throughout the book that symbols are in italic, while units of measurement are not. For example, P = F/A, with pascals Pa. Or *m* is mass while m is meter.

CONCEPT: For a constant force, **CHECK**: an increase in the area of contact will result in a decrease in the pressure.

Teaching Resources

- Reading and Study Workbook
- Problem-Solving Exercises
- in Physics 4-3
 PresentationEXPRESS
- Interactive Textbook





FIGURE 6.7 🔺

The author applies a force to fellow physics teacher Paul Robinson, who is bravely sandwiched between beds of sharp nails. The driving force per nail is not enough to per nail is not enough to puncture the skin. CAUTION: Do not attempt this on your own!



In attempting to do the demonstration shown in Figure 6.7, would it be wise to begin with a few nails and work upward to more nails? Answer: 6.5

> You exert more pressure against the ground when you stand on one foot than when you stand on both feet. This is due to the decreased area of contact. Stand on one toe like a ballerina and the pressure is huge. The smaller the area supporting a given force, the greater the pressure on that surface.

You can calculate the pressure you exert on the ground when you are standing. One way is to moisten the bottom of your foot with water and step on a clean sheet of graph paper. Count the number of squares on the graph paper contained within your footprint. Divide your weight by this area and you have the average pressure you exert on the ground when standing on one foot. How will this pressure compare with the pressure you exert when you stand on two feet?

A dramatic illustration of pressure is shown in Figure 6.7. The author applies appreciable force when he breaks the cement block with the sledgehammer. Yet his friend (the author of the lab manual) sandwiched between two beds of sharp nails is unharmed. The friend is unharmed because much of the force is distributed over the more than 200 nails that make contact with his body. The combined surface area of this many nails results in a tolerable pressure that does not puncture the skin. **CAUTION:** *This demonstration is quite dangerous. Do not attempt it on your own*.

CONCEPT How does the area of contact affect the pressure a **CHECK** force exerts on an object?

6.6 Free Fall Explained

Recall that free fall occurs when a falling object encounters no air resistance. Also recall that Galileo showed that falling objects accelerate equally, regardless of their masses. This is *strictly* true if air resistance is negligible, that is, if the objects are in free fall. It is *approximately* true when air resistance is very small compared with the mass of the falling object. For example, a 10-kg cannonball and a 1-kg stone dropped



from an elevated position at the same time will fall together and strike the ground at practically the same time. This experiment, said to be done by Galileo from the Leaning Tower of Pisa and shown in Figure 6.8, demolished the Aristotelian idea that an object that weighs ten times as much as another should fall ten times faster than the lighter object. Galileo's experiment and many others that showed the same result were convincing. But Galileo couldn't say *why* the accelerations were equal. The explanation is a straightforward application of Newton's second law and is the topic of the cartoon "Backyard Physics." Let's treat it separately here.

Recall that mass (a quantity of matter) and weight (the force due to gravity) are proportional. A 2-kg bag of nails weighs twice as much as a 1-kg bag of nails. So a 10-kg cannonball experiences 10 times as much gravitational force (weight) as a 1-kg stone. The followers of Aristotle believed that the cannonball should accelerate at a rate ten times that of the stone, because they considered only the cannonball's ten-times-greater weight. However, Newton's second law tells us to consider the mass as well. A little thought will show that ten times as much force acting on ten times as much mass produces the same acceleration,

$$\frac{F}{m} = \frac{F}{m}$$

where F stands for the force (weight) acting on the cannonball, and m stands for the correspondingly large mass of the cannonball. The small F and m stand for the smaller weight and smaller mass of the stone. As Figure 6.9 shows, the *ratio* of weight to mass is the same for these or any objects. All freely falling objects undergo the same acceleration at the same place on Earth. In Chapter 4 we introduced the symbol g for the acceleration.

FIGURE 6.8

In Galileo's famous demonstration, a 10-kg cannonball and a 1-kg stone strike the ground at practically the same time.

Remember that only a single force acts on something in free fall the force due to gravity.





FIGURE 6.9 🔺

The ratio of weight (F) to mass (m) is the same for the 10-kg cannonball and the 1-kg stone.

6.6 Free Fall Explained

Common Misconception Heavy objects always fall faster than

Ight objects. FACT This is only true in the presence of air resistance.

Demonstration

Drop a book and a sheet of paper to show the different falling rates. Then crumple up the piece of paper into a ball and show that it and the book accelerate almost equally when dropped.

Teaching Tip State that air resistance for the low speeds involved in the above demonstration does not reveal itself in your demonstration. Do not discuss the effects of air resistance until you have first investigated the physics that occurs in the absence of air resistance.

Teaching Tip Emphasize that free fall means falling free of air resistance or other constraints. Teach the physics of free fall first, and then consider nonfree fall.

A too-early preoccupation with air resistance can hide some very basic physics. Your selfdiscipline may be challenged by overanxious students who ask questions about air drag. Defer such questions until the simplest case is understood by your class.



NEWTON'S SECOND LAW OF MOTION—FORCE AND ACCELERATION 93

Teaching Tip State that Galileo, who is reputed to be the first to publicly show equal accelerations for unequal masses, could not adequately explain why. He lacked the model offered by Newton, namely Newton's second law.

Demonstration

either. Bring both arguments each argument seemed to with the expression " $a \sim F$," and so will take more time to a good argument for the cannonball and stone in this to the case of the falling second law, *a* = *Flm*. Relate together with Newton's be, the results do not support show that however reasonable and the chalk together to "a ~ 1/m." Drop the 1-kg mass and the second argument with Summarize the first argument get moving than the chalk.) more inertia than the chalk, argument for the chalk falling chalk.) Then articulate a good more by gravity than the (e.g., the 1-kg mass is pulled heavy mass falling faster. exposed to this idea. Articulate imagine they have not been the ground first if you release Ask the class which will hit of chalk above your head. Hold a 1-kg mass and a piece faster. (e.g., the 1-kg mass has "the same.") Ask students to (Your class should answer, them at the same time.



94

Figure 6.9.

object is the force due to gravity—the object's weight. Using Newton's cannonball is 100 N at Earth's surface. The force acting on a falling of a 1-kg stone is 10 N at Earth's surface. The weight of a 10-kg second law, the acceleration of the stone is We can show the same result with numerical values. The weight

$$= \frac{F}{m} = \frac{\text{weight}}{m} = \frac{10 \text{ N}}{1 \text{ kg}} = \frac{10 \text{ kg} \cdot \text{m/s}^2}{1 \text{ kg}} = 10 \text{ m/s}^2 = g$$

and the acceleration of the cannonball is

$$a = \frac{F}{m} = \frac{\text{weight}}{m} = \frac{100 \text{ N}}{10 \text{ kg}} = \frac{100 \text{ kg} \cdot \text{m/s}^2}{10 \text{ kg}} = 10 \text{ m/s}^2 = g$$

acceleration because the net force on an object is only its weight, discussed. Now we know why the acceleration of the coin and the and the ratio of weight to mass is the same for all objects. discussed in Chapter 4, the reason for the equal accelerations was not feather are the same. 🕑 All freely falling objects fall with the same In the famous coin-and-feather-in-a-vacuum-tube demonstration

CHECK: same acceleration? **CONCEPT** Why do all freely falling objects fall with the

6.7 Falling and Air Resistance

very unequally in the presence of air. When falling in air, the coin air resistance diminishes the net force acting on the falling objects. falls quickly while the feather flutters to the ground. The force due to The feather and coin fall with equal accelerations in a vacuum, but

describes the relationship between speed, area, and air resistance: experiences depends on the object's speed and area. An expression find that the force of air resistance is also proportional to the fronward, exposing maximum frontal area for the book, the air resistance hold your physics book out the window with the large side facing forresistance force depends on speed. If instead of just your hand, you car moves faster, the force on your hand increases, indicating that air when you stick your hand out of the window of a moving car. If the Speed and Area The force due to air resistance is experienced tal area of the moving object. **W** The air resistance force an object force is much larger than it was on your hand at the same speed. You

Air resistance force \sim speed \times frontal area

portional to the speed and frontal area of an object. The expression shows that the air resistance force is directly pro-

> gravity and air resistance it is not in free fall. act on a falling object, When the forces of





Web Code: csn - 0607

Which experiences a elephant? of paper or a falling greater air resistance Answer: 6.7.1 force, a falling piece

> weight, and the ratio of weight same acceleration because the **CHECK** All freely falling to mass is the same for all objects net force on an object is only its

Teaching Resources

- Transparency 9
- Next-Time Questions 6-3, 6-4

Falling and Air Resistance

Key Terms

terminal speed, terminal velocity

Objects have no weight in a vacuum Common Misconceptions

object since a falling object in a effect on the weight of an object vacuum does not experience air and acceleration of a falling FACT The absence of air has no resistance. However, it does affect the speed

same speed as its firing speed. handgun normally returns with the Because of air resistance, a bullet tired vertically from a rifle or

bullet fired vertically from a gun FACT Because of air resistance, a than 10% its firing speed. normally returns at a speed less

objects, it is useful to exaggerate Similarly, but not as much, for the feather than for the falling rock. significant role for the falling the air resistance plays a more for the light person, and the substitute the falling of a feather skydiver, ask your students to the falls of a heavy and a light effects are more clearly visualized the circumstance so that the effects of air resistance on falling falls of the two skydivers. heavy person. It is easy to see that falling of a heavy rock for the For example, in comparing Teaching Tip In discussing the

Teaching Tip Discuss Figure 6.10 and Figure 6.11.

the weight of the falling body. depends on two things: (1) how there may be a point where the object falls faster and faster, on both size and speed. As an it's going. Air resistance depends disturbed air is and (2) how fast big it is, or how big the path of that a falling body encounters State that the air resistance torce of air resistance is equal to

does not mean zero velocity, but velocity)? No, zero acceleration zero acceleration means zero comes to an abrupt halt (i.e., that this mean the falling object acceleration then be? Zero Does the object? Zero What will the what will be the net force on equal to the object's weight, resistance on a falling object is Ask When the force of air

only difference will be that the second law will still be the mode net force is not simply the weight presence of air resistance. The zero CHANGE in velocity. for investigating falling in the Teaching Tip Newton's

object's weight. air resistance builds up to counter a falling object will decrease as or describe how the net force on acceleration will always be less of a falling object is $a = F_{net}/m =$ of the falling object, but the the accelerating effect of the class discussion to this equation, than g by the amount R/m. Direct that with air resistance the air resistance R, the acceleration weight minus air resistance. With (mg - R)/m = g - (R/m). Note (weight – resistance)/m =

velocity doubles when the mass is area. Therefore the terminal object divided by its surface mass is increased by nine times. quadrupled. The terminal velocity square root of the mass of an velocity is proportional to the increases by three times if the Teaching Tip The termina

> when air resistance equals weight. Sky divers reach terminal speed FIGURE 6.10

nor changing direction. up nor slowing down have, neither speeding eration means that the velocity. Zero acceltion does not mean zero velocity it happens to object will maintain the size that zero accelera-It's important to empha-





FIGURE 6.11

its fall. and decreases the speed of its area by spreading out. The flying squirrel increases This increases air resistance



objects, we say it has reached its terminal velocity. Terminal velocity weight. If we are concerned with direction, which is down for falling is terminal speed together with the direction of motion. acceleration of a falling object is zero because friction balances the reached its terminal speed. Terminal speed is the speed at which the is zero and the object stops accelerating. We say that the object has it equals the weight of the object, then the net force on the object like the sky divers shown in Figure 6.10, builds up to the point where **Terminal Speed** When the air resistance force on a falling object,

have to fall faster than a feather to reach its terminal speed. has a relatively small area compared to its weight, so the coin will resistance has a large effect on the feather's motion. A coin, however, is large relative to its very small weight. Even at small speeds the air A falling feather reaches its terminal speed quite quickly. Its area

son. The greater weight is more effective in "plowing through" air. the flying squirrel in Figure 6.11. heavier person will attain a greater terminal speed than a lighter per-200 km/h, depending on the weight and orientation of the body. A when the body is spread out and surface area is increased, like that of Body orientation also makes a difference. More air is encountered The terminal speed for a sky diver varies from about 150 to



to LIFE SCIENCE

snake (Chysopelea paradisi) jumps from a tree branch it doubles its increasing their surface areas when falling. When the paradise tree Terminal Velocity Skydivers and flying squirrels are not alone in than 20 meters in a single leap. maneuvers itself by undulating in a graceful S-shape, traveling more width by flattening itself. It acquires a slightly concave shape and

discover!

How Does Air Resistance Affect the Motion of Falling Objects?

- 1. Drop a sheet of paper and a book side-by-side at the same at the same time? time. Does either of them hit the ground first, or do they land
- 2. Place the piece of paper against the bottom surface of the hor of them hit the ground first, or do they land at the same time? izontally held book. Drop them at the same time. Does either
- **3.** Repeat Step 2 with the piece of paper on top of the book.
- 4. Think Explain the effect of air resistance on the motion of the piece of paper and the book in Steps 1–3.

squirrel while the light person falls head or feet first. A parachute 15 to 25 km/h, slow enough for a safe landing. greatly increases air resistance, and cuts the terminal speed down to proximity to each other if the heavy person spreads out like a flying tation. A heavy sky diver and a light sky diver can remain in close Terminal speed can be controlled by variations in body orien-

acceleration falling in air than a foam ball. than the baseball does. Figure 6.12 shows that a golf ball has a greater less for the tennis ball. The tennis ball behaves more like a parachute nis ball than on the heavier baseball, so the acceleration of the fall is is often negligible, but at high speeds, it can make quite a difference. buildup of air resistance at higher speeds. At low speeds, air resistance the heavier baseball strikes the ground first. This is due to the time. But if you drop them from the top of a building, you'll notice them at the same time, you'll see them strike the floor at the same The effect of air resistance is more pronounced on the lighter ten-If you hold a baseball and tennis ball at arm's length and release

stood until Newton announced his second law of motion. of Aristotle. The behavior of falling objects was never really underrather than the pronounced time difference expected by the followers ground first. However, the time difference was only a split second, from the Leaning Tower of Pisa, the heavier object did get to the When Galileo reportedly dropped the objects of different weights

primarily responsible for this feat. to men landing on the moon in the 1900s. Newton's second law was acceleration, force, and mass, discovered by Newton in the 1600s, led ing how concepts connect to one another. The connection between Isaac Newton truly changed our way of seeing the world by show-

CHECK : an object? **CONCEPT**: What factors determine the air resistance force on



FIGURE 6.12

a golf ball and a foam ball acceleration is greater. coming air resistance, so its ball is more effective in overfalling in air. The heavier golf This stroboscopic photo shows

think

each wears the same size the same altitude and parachutes together at light person open their Answer: 6.7.2 the ground first? parachute, who will reach If a heavy person and a

discover!

MATERIALS book, paper

and the book hit the ground In Steps 2 and 3, the paper the book hits the ground first at the same time. EXPECTED OUTCOME In Step 1,

you drop them, the book the book and the paper weight relative to the air faster because of its greater for the paper. below it and "clears the way" overcomes the air resistance before you drop them. When virtually no air between falls. In Step 3, there is pushes the paper with it as it resistance. In Step 2, the book THINK In Step 1, the book falls

because the net force acting will reach zero and she will have decrease, or remain the same? does her acceleration increase air resistance. As air resistance on her is her weight minus on her increase, decrease, or decrease, or remain the same? air, does air resistance increase, falls faster and faster through the a high-altitude balloon. As she Ask A skydiver jumps from net force decreases. When she Acceleration decreases because As she falls faster and faster, increases, net force decreases. remain the same? Decrease, Increase Does the net force reached terminal velocity. falls fast enough, her acceleration

experiences depends on the object's speed and area. **CHECK** force an object

 Next-Time Questions 6-5, 6-6, 6-7

Teaching Resources

CHAPTER 6

NEWTON'S SECOND LAW OF MOTION—FORCE AND ACCELERATION 97



cause motion. force changes motion. It doesn't Teaching Tip Remember that

Teaching Tip An important act directly on the object. draw a force vector diagram that Newton's second law is to first step in solving problems with includes only those forces that



Go Vnline PHSchool.com Web Code: csa – 0600 Visit: PHSchool.com For: Self-Assessment

Concept Summary

- Unbalanced forces acting on an object cause the object to accelerate.
- e mass will result in a decrease in the For a constant force, an increase in the acceleration.
- direction as the net force, and is inversely magnitude of the net force, is in the same an object is directly proportional to the celeration produced by a net force on Newton's second law states that the acproportional to the mass of the object.
- contact and how much the surfaces are pressed together. faces depends on the kinds of material in The force of friction between two sur-
- area of contact will result in a decrease in For a constant force, an increase in the the pressure.
- e All freely falling objects fall with the same weight to mass is the same for all objects object is only its weight, and the ratio of acceleration because the net force on an
- pends on the object's speed and area. The air resistance force on an object de-

Key Terms

ree-body diagram (p. 91)	luid (p. 90)	second law (p. 88)	nversely (p. 87) Jewton's
(p. 96)	(p. 96) terminal velocity	terminal speed	pressure (p. 91)

think! Answers

- 6:3 The same force on twice the mass produces half the acceleration, or 1 m/s²
- 6.4 No, not unless the book tends to slide or is sliding. (More about this in the Conceptoccur only when an object tends to slide or will act toward the right. Friction forces then friction between the book and table is pushed toward the left by another force, does slide across the table. For example, if it Development Practice Book.)
- 6.5 No, no, no! There would be one less physics sure would cause harm. with fewer nails. The resulting greater presteacher if the demonstration were performed
- 6.7.1 The elephant! It has a greater frontal area story! molecules out of the way. The effect of the air of which mean the elephant pushes more air and falls faster than a piece of paper-both resistance force on each, however, is another
- 6.7.2 The heavy person will reach the ground first speed is reached. The heavy person moves ahead of the light person, and the separation continues to accelerate until a greater terminal minal speed sooner, while the heavy person Like a feather, the light person reaches tercontinues to increase as they descend

86

• Virtual Physics Labs 8, 9 • Conceptual Physics Alive! DVDs Newton's Second Law

TeacherEXPRESS

Teaching Resources



Check Concepts

Section 6.1

1. What produces acceleration?

2. In Chapter 4 we defined acceleration as the time rate of change of velocity. What other equation for acceleration is given in this chapter?

Section 6.2

3. Is acceleration directly proportional to mass, or is it inversely proportional to mass?



4. If two quantities are inversely proportional to each other, does that mean as one increases the other increases also?

Section 6.3

- **5.** If the net force acting on a sliding block is tripled, what happens to the acceleration?
- **6.** If the mass of a sliding block is tripled at the same time the net force on it is tripled, how does the resulting acceleration compare with the original acceleration?

Section 6.4

7. Motion is affected by solid objects in contact. In what other situations does friction affect motion?

- 8. Suppose you exert a horizontal push on a crate that rests on a level floor, and it doesn't move. How much friction acts compared with your push?
- 9. How great is the air resistance that acts on a 10-N sack that falls in air at constant velocity?

Section 6.5

10. Distinguish between force and pressure.

11. When do you produce more pressure on the ground, standing or lying down?



12. Why is it important that many nails are in the boards of Figure 6.7?

Section 6.6

13. What is meant by *free fall*?

- 14. The ratio of circumference/diameter for all circles is π . What is the ratio of force/mass for all freely-falling bodies?
- 15. Why doesn't a heavy object accelerate more than a light object when both are freely falling?

ASSESS

Check Concepts

- **1.** Net force
- **2.** *a* = *F*/*m*
- **3.** Acceleration is inversely proportional to mass.
- Increasing mass means decreasing acceleration, for example.
- **4.** No, just the opposite
- 5. Acceleration also triples.
- **6.** Triple the force on three
- times the mass results in no change in acceleration.
- In fluids, through a liquid or air
- 8. Same magnitude as your push, oppositely directed, producing a zero net force
- 9. 10 N; then the net force equals zero.
- Force is a push or pull; pressure is force per unit area.
- **11.** More pressure when you're standing due to less area of contact for the same force
- 12. More nails, more surface area of nails against his body, so less pressure when hammer hits
- Free fall occurs when only the force of gravity acts; no air resistance. (Different meaning to a sky diver, where "free fall" means falling in air before a parachute is opened.)
 The ratio is or the acceleration
- **14.** The ratio is *g*, the acceleration of free fall.
- **15.** Greater mass means proportionally greater weight and, hence, proportionally more force. The ratio force/ mass is unchanged with equal

acceleration in free fall.

- Increases; more air/second is plowed through as speed increases
- increases. **17.** The faster falling object
- 18. Zero
- **19.** Surface area, depending on body orientation; more area to the air, more resistance
- **20.** 100 N, for then the net force will be zero

Think and Rank

21. D, B, A, C
22. a. D, A = B = C
b. A = C, B = D
23. a. B, C, A
b. C, A = B

24. C, D, B, A

ASSESS (continued)

Section 6.7

- **16.** Does air resistance on a falling object increase or does it decrease with increasing speed?
- **17.** If two objects of the same size fall through air at different speeds, which encounters the greater air resistance?
- **18.** What is the acceleration of a falling object that has reached its terminal velocity?
- **19.** What, besides speed, affects the air resistance on a skydiver?
- 20. How much air resistance acts on a falling 100-N box of nails when it reaches terminal velocity?

Think and Rank

Rank each of the following sets of scenarios in order of the quantity or property involved. List them from left to right. If scenarios have equal rankings, then separate them with an equal sign. (e.g., A = B)

21. Each diagram shows a ball traveling from left to right. The position of the ball each second is indicated by the second. Rank the net forces from greatest to least required to produce the motion indicated in each diagram. Right is positive and left is negative.



22. Boxes of various masses are on a friction-free level table.



Rank each of the following from greatest to least.

- **a.** the net forces on the boxes
- **b.** the accelerations of the boxes
- 23. Each block on the friction-free lab bench is connected by a string and pulled by a second falling block.



Rank each of the following from greatest to least.

- a. the acceleration of the two-block systemsb. the tension in the strings
- 24. All the aluminum blocks have the same mass and are gently lowered onto a gelatin surface, which easily supports them. All have square bottom surfaces. Rank them by how much they dent into the surface, from greatest to least depth.



gelatin surface

100

Plug and Chug

These questions are to familiarize you with the key equations of the chapter.

Acceleration = $\frac{\text{net force}}{\text{mass}}$

 $a = \frac{F}{m}$

- **25.** Calculate the acceleration of a 40-kg crate of softball gear when pulled sideways with a net force of 200 N.
- **26.** Calculate the acceleration of a 2000-kg, single-engine airplane as it begins its takeoff with an engine thrust of 500 N.
- **27.** Calculate the acceleration of a 300,000-kg jumbo jet just before takeoff when the thrust for each of its four engines is 30,000 N.
- **28.** Calculate the acceleration if you push with a 20-N horizontal force against a 2-kg block on a horizontal friction-free air table.

F = ma

- **29.** Calculate the horizontal force that must be applied to a 1-kg puck to make it accelerate on a horizontal friction-free air table with the same acceleration it would have if it were dropped and fell freely.
- **30.** Calculate the horizontal force that must be applied to produce an acceleration of 1.8g for a 1.2-kg puck on a horizontal friction-free air table.

Think and Explain

- **31.** If you push horizontally on your book with a force of 1 N to make the book slide at constant velocity, how much is the force of friction on the book?
- **32.** Terry says that if an object has no acceleration, then no forces are exerted on it. Sherry doesn't agree, but can't provide an explanation. They both look to you. What do you say?
- **33.** When a car is moving in reverse, backing from a driveway, the driver applies the brakes. In what direction is the car's acceleration?
- **34.** The auto in the sketch moves forward as the brakes are applied. A bystander says that during the interval of braking, the auto's velocity and acceleration are in opposite directions. Do you agree or disagree?



35. What is the difference between saying that one quantity is proportional to another and saying it is equal to another?

Plug and Chug.....

25. a = F/m = (200 N)/(40 kg) = 5 m/s²

26. *a* = *F*/*m* = (500 N)/(2000 kg) = 0.25 m/s²

27. *a* = *F/m* = 4(30,000 N)/ (300,000 kg) = 0.4 m/s²

28. *a* = *F/m* = (20 N)/(2 kg) = 10 m/s²

29. *F* = *ma* = (1 kg)(10 m/s²) = 10 N

30. F = ma = (1.2 kg)(1.8)(10 m/s²) = 21.6 N

Think and Explain

- **31.** 1 N in the opposite direction to the push
- **32.** Say that Terry would be correct if she said "net" force.
- **33.** When the car moves backward, acceleration is forward in direction of the net force on it, decreasing the speed of the car.
- **34.** Agree, for acceleration is opposite his velocity. (He is decelerating.)
- 35. Proportional—a change in one quantity implies a corresponding change in the other; Identical—when one quantity is equal to another

- **36.** Always g; $a = F/m = mg/m = g \neq 0$
- **37.** To reduce ground contact and friction between the opponent's feet and the ground
- **38.** At the beginning of takeoff, when air drag is least
- 39. By a = F/m, as m decreases, a increases for a given force. Rocket's mass becomes less as fuel is burned. (Also, the gravitational force is less with distance from Earth.)
- **40.** By *a* = *F/m*, a sudden stop means a great *a*, which means a great *F* for a given mass.
- **41.** On the middle hill, where the slope becomes less toward the bottom
- **42.** Smaller cutting area produces a greater pressure for a given force.
- **43.** No, the scale reads force, not pressure.
- **44.** By a = F/m, F equals mg in free fall, a = mg/m = g. Mass cancels out.
- **45.** Both are correct. The effects of force/inertia are not confined to free fall.



- **36.** What is the acceleration of a rock at the top of its trajectory when thrown straight upward? Explain whether or not the answer is zero by using the equation a = F/m as a guide to your thinking.
- **37.** When blocking in football, why does a defending lineman often attempt to get his body under that of his opponent and push upward? What effect does this have on the friction force between the opposing lineman's feet and the ground?



- **38.** An aircraft gains speed during takeoff due to the constant thrust of its engines. When is the acceleration during takeoff greatest—at the beginning of the run along the runway or just before the aircraft lifts into the air? Think, then explain.
- **39.** A rocket becomes progressively easier to accelerate as it travels through outer space. Why is this so? (*Hint:* About 90 percent of the mass of a newly launched rocket is fuel.)



- **40.** A common saying goes, "It's not the fall that hurts you; it's the sudden stop." Translate this into Newton's laws of motion.
- **41.** On which of these hills does the ball roll down with increasing speed and decreasing acceleration along the path? (Use this example if you wish to explain to someone the difference between speed and acceleration.)



- **42.** Why does a sharp knife cut better than a dull knife?
- **43.** When Helen lifts one foot and remains standing on a bathroom scale, pressure on the scale is doubled. Does the weight reading change?
- **44.** Aristotle claimed the speed of a falling object depends on its weight. We now know that objects in free fall, whatever their weights, gain speed at the same rate. Why does weight not affect acceleration?
- **45.** After learning why objects of different mass have the same acceleration in free fall, Erik wonders if objects tied to equal lengths of string would swing together in unison. Lisa wonders if objects of different masses would slide at equal speeds down a friction-free inclined plane. What is your thinking on these hypotheses?

- **46.** In a vacuum, a coin and a feather fall side by side. Would it be correct to say that in a vacuum equal forces of gravity act on both the coin and the feather?
- **47.** As a sky diver falls faster and faster through the air (before reaching terminal speed), does the net force on her increase, decrease, or remain unchanged? Does her acceleration increase, decrease, or remain unchanged? Defend your answers.
- **48.** After she jumps, a sky diver reaches terminal speed after 10 seconds. Does she gain more speed during the first second of fall or the ninth second of fall? Compared with the first second of fall, does she fall a greater or a lesser distance during the ninth second?
- **49.** Can you think of a reason why the acceleration of an object thrown downward through the air would actually be less than 10 m/s²?
- **50.** How does the weight of a falling body compare with the air resistance it encounters just before it reaches terminal velocity? Just after it reaches terminal velocity?
- **51.** Why does a cat that falls from a 50-story building hit the safety net with no more speed than if it fell from the 20th story?



- **52.** A regular tennis ball and another one filled with sand are dropped at the same time from the top of a high building. Your friend says that even though air resistance is present, both balls should hit the ground at the same time because they are the same size and pass through the same amount of air. What do you say?
- **53.** If you drop an object, its acceleration toward the ground is 10 m/s². If you throw it downward instead, will its acceleration after throwing be greater than 10 m/s²? Why or why not? (Ignore air resistance.)
- **54.** Suzy Skydiver, who has mass *m*, steps from the basket of a high-flying balloon of mass *M* and does a sky dive.
- **a.** What is the net force on Suzy at the moment she steps from the basket?
- **b.** What is the net force on her when air resistance builds up to equal *half* her weight?
- **c.** What is the net force on her when she reaches terminal speed *v*?
- **d.** What is the net force on her after she opens her parachute and reaches a new terminal speed 0.1*v*?

Think and Solve

55. What is the acceleration during takeoff of a jumbo jet with a mass of 30,000 kg when the thrust for each of its *four* engines is 30,000 N?

- 46. No, no! The reason is that ratios of weight/mass are equal.
- 47. Decreases; decreases; resisted by air drag
- 48. Maximum gain in speed occurs during the first second of fall. Since her speed increases, she will be falling faster during the last second of fall, and therefore cover the greatest distance during
- **49.** With air drag, acceleration is actually less for higher speed.

this time.

- **50.** Just before terminal velocity, weight is slightly greater than air drag. After reaching terminal velocity, weight is equal and opposite to air drag.
- Once anything reaches terminal velocity, no increase in speed occurs. This occurs for a cat at about the 20th story.
- **52.** The heavier ball falls through the air with greater acceleration, so it hits the ground first.
- 53. The speed is different but the acceleration is still 10 m/s². It still gains 10 m/s for each second.
- **54.** a. With no air drag yet, $F_{net} = mg$ (her weight).
- **b.** $F_{net} = mg R = mg (mg/2) = mg/2$ (half her weight)
- **c.** At terminal velocity, a = 0 so the net force = 0.
- **d**. At any terminal velocity, net force = 0.

Think and Solve

55. $a = F/m = (4 \times 30,000 \text{ N}) \div$ (30,000 kg) = 4 m/s²

67. a. $a = F_{net}/m = (100 \text{ N} - 100 \text{ N})$ **66.** $a = (1 \text{ kg} \times g)/(1 \text{ kg} + 100 \text{ kg})$ **64.** $a = F_{net}/m = (mg - R)/m$, so **61.** If *Flm* = 1 m/s², then **60.** $a = F_{net}/m$ so F = ma**59.** $a = F_{net}/m$ so m = F/a**58.** $a = \Delta v / \Delta t = (0 \text{ m/s} - 9 \text{ m/s}) \div$ **57.** *a* = *F*_{net}/*m* = 10.0 N/ 6.7 kg 56. Half as much, 1.5 m/s²; same **65.** $F = ma = 10 \text{ kg} (10 \text{ m/s}^2) =$ **63.** $P = F/A = (20 \text{ N})/(0.05 \text{ m}^2)$ **62.** *F* = *ma* 5 m/s≁ 20 N / $25 \text{ kg} = 3.2 \text{ m/s}^2$ maximum acceleration = gso a is negative. Also R > mgR = mg - ma = m(g - a) =50 kg [10m/s² - (-6.2 m/s²)] $(20 \text{ N})/(0.01 \text{ m}^2) = 2000 \text{ N/m}^2$ = 21,000 N (about 4620 lb) = 70 kg \times 30 \times 10 m/s² (Note 10 N = 10 kg·m/s².) = 10.0 N/2.0 m/s² = 5.0 kg $0.2 s = -45 m/s^2$ $= 1.5 \text{ m/s}^2$ **b.** $d = 1/2at^2 =$ = 810 N. Note a is opposite g F/(0.75m) = 1.33 m/s² = m ($\Delta v/\Delta t$) = 2000 kg(0 m/s force acts on twice the mass (100/101)g = 0.99g;(100 kg + 1 kg) = $= (1/101)g; a = (100 \text{ kg} \times g)/$ 100 N; a = (100 N)/(20 kg) =for a net upward force. = 400 N/m²; P = F/A = $1/2(3.2 \text{ m/s}^2) (5 \text{ s})^2 = 40 \text{ m}$ 22 m/s/4.0 s = -11,000 N

Activities

68. Air resistance slows the brief fall, but from a higher might not be noticed for a the coin still falls faster. This so the coin hits the ground more than the dropped coin, dropped sheet of paper much first. Crumpled paper will fall fall, the coin decidedly wins frontal area to the air, but faster because it presents less



- 56. A net force on a 2-kg cart accelerates the the same net force produce on a 4-kg cart? cart at 3 m/s². How much acceleration will
- **57.** A net force of 10.0 N is exerted by Irene the cart will have an acceleration of 1.5 m/s^2 . on a 6.7-kg cart for 3.0 seconds. Show that
- 58. Toby Toobad, who has a mass of 100 kg, is skateboarding at 9.0 m/s when he smacks (that's 4.5 times g-ouch!). 0.2 s. Show that his deceleration is 45 m/s^2 into a brick wall and comes to a dead stop in
- 59. A net force of 10.0 N on a box of plastic Show that the mass of the box is 5.0 kg. toam causes it to accelerate at 2.0 m/s^2 .



- **60.** Austin's truck has a mass of 2000 kg. When ing force acting on the truck is 11,000 N. 4.0 s. Show that the magnitude of the braktraveling at 22.0 m/s, it brakes to a stop in
- **61.** If a loaded truck that can accelerate at 1 m/s^2 if the same driving force acts on it? original mass, what acceleration can it attain loses its load and has three-fourths of its
- 62. An occupant of a car can survive a crash if decelerating at this rate. 30g. Calculate the force on a 70-kg person the deceleration during the crash is less than

- **63.** What is the pressure on a table when a 20-N book with a 0.05-m² cover lies flat on it? on its end (area 0.01 m²)? What is the pressure when the book stands
- 64. A falling 50-kg parachutist experiences an force is 810 N when this occurs. opens her parachute. Show that the drag upward acceleration of 6.2 m/s² when she
- 65. A 10-kg mass on a horizontal friction-free system of both masses? due to gravity, in newtons, on the hanging to another 10-kg mass hanging vertically air track is accelerated by a string attached from a pulley as shown. What is the force 10-kg mass? What is the acceleration of the



66. Suppose the masses described in problem pare the accelerations when they are inter-65 are 1 kg and 100 kg, respectively. Commaximum acceleration of such a system of mass dangles over the pulley, and then for changed, that is, for the case where the 1-kg masses? the pulley. What does this indicate about the the case where the 100-kg mass dangles over

- 67. Skelly the skater is propelled by rocket power. Skelly and the rocket together have a mass of 25 kg. The thrusting force is 100 N and friction is 20 N.a. What is Skelly's acceleration?
- How far does he go in 5 s if he starts from rest?

<u>ь</u>



Activities

- **68.** Drop a sheet of paper and a coin at the same time. Which reaches the ground first? Why? Now crumple the paper into a small, tight wad and again drop it with the coin. Explain the difference observed. Will the coin and paper fall together if dropped from a second-, third-, or fourth-story window? Try it and explain your observations.
- **69.** Glue a penny to a string. When in a moving automobile, hang the string and penny out a window. It will be swept backward due to air resistance. When the string makes an angle of 45°, easily seen with a protractor, the air resistance on the coin equals the weight of the coin. A look at the speedometer tells you the coin's terminal speed in air! Do you see why the angle makes a difference?

- **70.** The net force acting on an object and the resulting acceleration are always in the same direction. You can demonstrate this with a spool. If the spool is gently pulled horizon-tally to the right, in which direction will it roll?
- 71. Write a letter to a friend who has not yet studied physics and tell what you've learned about Galileo introducing the concepts of acceleration and inertia. Tell of how Galileo was also familiar with forces, but didn't see the connection, revealed in his second law of motion. Explain with the second law why heavy and light objects in free fall gain the same speed in the same time. In this letter, it's okay to use an equation or two, making it clear that you see equations as a shorthand notation of explanations.



- **69.** A 45° angle merely simplifies the trigonometry. At 45° the magnitudes of the force of gravity and air drag on the coin are equal. Both forces make up the sides of a right triangle. At other angles, more trigonometry would be needed.
- **70.** If the string is pulled a supporting view of this of pull). Net force and without rolling-but still a critical angle (where the if you pull the string vertically, of pull and acceleration are Development Practice Book section of the Conceptto this activity in the torque direction. You will return acceleration are in the same the spool will be dragged contact with the surface), isn't perfectly vertical. At even do so if the string opposite direction. It will the spool will roll in the clearly the same. Interestingly, horizontally, the direction (and see how torque gives the horizontal component (actually in the direction of in the direction of the pull line of action of the string intercepts the point of wheel
- Answers will vary.

activity).



105

Computer Test Bank
Chapter and Unit Tests

Teaching Resources

-