Alg 2 Final Exam Review Probability and Statistics


Spring 2014

- 1. There are 12 different flags that you could put out front of the school building..
- a) If there are 12 flag poles out front how many different ways could you arrange all twelve flags on the poles?
- b) If three of the poles have broken ropes to hoist the flags, how many different ways could you arrange 9 of the flags?
- 2. You have 5 cans of paint in the basement and you want to mix some of them together to paint the basement walls...
- a) How many different new colors could you create if you mix all 5 of them together??
- b) How many different new colors could you create if you can only mix 3 of them together?
- 3. The push button lock on your car has a password that must be 5 characters long, 3 of the characters must be numbers (0 to 9) and 2 of the characters must be letters of the alphabet.:
- a) How many different passwords are possible if Letters and Numbers can't repeat?
- b) How many different passwords are possible if Numbers can't repeat but letters can?.
- 5. Use this set 10 scores: 23,24,33,52,57,98,99,103,104,122
- a) What number is at the 40th percentile?
- b) 99 is at what percentile?
- 8. Use the results of the survey shown below of what elementary students are afraid of. You will select one person at random. Find each probability as a fraction.

	Spiders	Darkness	Dogs	Snakes	Total
Boys	30	50	25	70	175
Girls	60	15	10	40	125
Totals	90	65	35	110	300

- a) P(boy or spiders)
- b) P(girl and snakes)

- c) P(Dogs or Darkness)
- d) P(snakes | boy)
- e) P(girl | spider)
- 9. You will spin the spinner once. Find each probability as a fraction.



- a) P(Red or Even)
- b) P(Purple or Green)

- c) P(Odd and Purple)
- d) P(Factor of 6 and Red)
- 10. In your Halloween bag of candy there are 5 Hershey bars, 6 Snickers, and 4 Reeses. In the morning you reach and grab a random piece of candy look at it, put it back into the bag, then grab another random piece of candy. Find each probability as a fraction. Do not reduce.
- a) P(Hershey bar and then Snickers)


b) P(Reeses and then another Reeses)

- 11. In your refrigerator you have the following drinks: 10 Pepsi's, 20 Cokes, and 3 Dr. Peppers. You take one at random and drink it. Realizing you are still thirsty you take another random drink and finish it. Find each probability as a fraction. Do not reduce.
- a) P(Coke and then another Coke)

- b) P(Dr. Pepper and then a Pepsi)
- 12. You go to the buffet for lunch and by a 3 item plate. You have 8 salads to choose from, 10 kinds of pasta, and 11 desserts. If you can only choose one of each how many different 3 item plates could you purchase?
- 13. The probability that I'll have a Coke at the ballgame is $\frac{7}{12}$ and the probability that I'll have a hot dog is $\frac{2}{9}$. Find the probability that I'll have a Coke or a hot dog at the game. Give your answer as a percent rounded to the nearest tenth.

P(Coke or hot dog)

- 14. At a party some door prizes are given away to the guests. Each guest will be given one door prize. The probability that you win tickets to a concert are $\frac{3}{11}$ and the probability that you win a free dinner are $\frac{2}{7}$. Find the probability that you win the tickets or the dinner as a percent rounded to the nearest tenth. P(Tickets or Dinner)
- 15. A survey of 900 people shows that 611 of them approve of the presidents job so far.
- a) Find the sample proportion to the nearest whole percent.
- b) Find the margin of error in the survey to the nearest whole percent.
- c) Find the interval that likely contains the actual population proportion.
- 16. A national test has a mean of 170 and a standard deviation of 8.
- a) Find the percent of tests that are below 186.
- b) Find the percent of tests that are above 162.
- c) Find the percent of data that are between 154 and 178.
- d) If a test is taken at random find the probability that the score is between 146 and 162.
- 17. Use the Venn Diagram below. Find each probability as a fraction without reducing.



- a) P(Coke but not Pepsi)
- b) P(7-Up or Coke)
- c) P(Pepsi and 7-Up)

- d) P(Not Pepsi)
- e) P(Coke or Pepsi but not 7-Up)